arXiv:1702.00561v1 [math.GR] 2 Feb 2017

# Autocommuting probability of a finite group

Parama Dutta and Rajat Kanti Nath*

*Department of Mathematical Sciences, Tezpur University, Napaam-784028, Sonitpur, Assam, India.*
*Emails: parama@gonitsora.com and rajatkantinath@yahoo.com*

**Abstract**

Let $G$ be a finite group and $\mathrm{Aut}(G)$ the automorphism group of $G$. The autocommuting probability of $G$, denoted by $\Pr(G, \mathrm{Aut}(G))$, is the probability that a randomly chosen automorphism of $G$ fixes a randomly chosen element of $G$. In this paper, we study $\Pr(G, \mathrm{Aut}(G))$ through a generalization. We obtain a computing formula, several bounds and characterizations of $G$ through $\Pr(G, \mathrm{Aut}(G))$. We conclude the paper by showing that the generalized autocommuting probability of $G$ remains unchanged under autoisoclinism.

*Key words:* Automorphism group, Autocommuting probability, Autoisoclinism.
*2010 Mathematics Subject Classification:* 20D60, 20P05, 20F28.

## 1 Introduction

Let $G$ be a finite group and $\mathrm{Aut}(G)$ be its automorphism group. For any $x \in G$ and $\alpha \in \mathrm{Aut}(G)$ the element $x^{-1}\alpha(x)$, denoted by $[x, \alpha]$, is called an autocommutator in $G$. We write $S(G, \mathrm{Aut}(G))$ to denote the set $\{[x, \alpha] : x \in G \text{ and } \alpha \in \mathrm{Aut}(G)\}$ and $K(G) := \langle S(G, \mathrm{Aut}(G))\rangle$. We also write $L(G) := \{x : [x, \alpha] = 1 \text{ for all } \alpha \in \mathrm{Aut}(G)\}$. Note that $K(G)$ and $L(G)$ are characteristic subgroups of $G$ known as the autocommutator subgroup and absolute center of $G$. It is easy to see that $K(G)$ contains the commutator subgroup $G'$ and $L(G)$ is contained in the center $Z(G)$ of $G$. Further, $L(G) = \bigcap\limits_{\alpha \in \mathrm{Aut}(G)} C_G(\alpha)$, where $C_G(\alpha) = \{x \in G : [x, \alpha] = 1\}$ is a subgroup of $G$ known as acentralizer of $\alpha \in \mathrm{Aut}(G)$. The subgroups $K(G)$ and $L(G)$ were defined and studied by Hegarty in [6]. Let $C_{\mathrm{Aut}(G)}(x) := \{\alpha \in \mathrm{Aut}(G) : \alpha(x) = x\}$ for $x \in G$ then $C_{\mathrm{Aut}(G)}(x)$ is a subgroup of $\mathrm{Aut}(G)$ and $C_{\mathrm{Aut}(G)}(G) = \{\alpha \in \mathrm{Aut}(G) : \alpha(x) = x \text{ for all } x \in G\}$. It follows that $C_{\mathrm{Aut}(G)}(G) = \bigcap\limits_{x \in G} C_{\mathrm{Aut}(G)}(x)$.

*Corresponding author

Let $G$ be a finite group acting on a set $\Omega$. In the year 1975, Sherman [13] introduced the probability that a randomly chosen element of $\Omega$ fixes a randomly chosen element of $G$. We denote this probability by $\Pr(G, \Omega)$. If $\Omega = \mathrm{Aut}(G)$ then $\Pr(G, \mathrm{Aut}(G))$ is nothing but the probability that the autocommutator of a randomly chosen pair of elements, one from $G$ and the other from $\mathrm{Aut}(G)$, is equal to 1 (the identity element of $G$). Thus

$$\Pr(G, \mathrm{Aut}(G)) = \frac{|\{(x, \alpha) \in G \times \mathrm{Aut}(G) : [x, \alpha] = 1\}|}{|G||\mathrm{Aut}(G)|}. \tag{1.1}$$

$\Pr(G, \mathrm{Aut}(G))$ is called autocommuting probability of $G$. Sherman [13] considered the case when $G$ is abelian and $\Omega = \mathrm{Aut}(G)$. Note that if we take $\Omega = \mathrm{Inn}(G)$, the inner automorphim group of $G$, then $\Pr(G, \mathrm{Inn}(G))$ gives the probability that a randomly chosen pair of elements of $G$ commute. $\Pr(G, \mathrm{Inn}(G))$ is known as commuting probability of $G$. It is also denoted by $\Pr(G)$. The study of $\Pr(G)$ was initiated by Erdös and Turán [4]. After Erdös and Turán many authors have worked on $\Pr(G)$ and its generalizations (conf. [3] and the references therein). Somehow the study of $\Pr(G, \mathrm{Aut}(G))$ was neglected for long time. At this moment we have only a handful of papers on $\Pr(G, \mathrm{Aut}(G))$ (see [2], [8], [12]). In this paper, we study $\Pr(G, \mathrm{Aut}(G))$ through a generalization.

In the year 2008, Pournaki and Sobhani [10] have generalized $\Pr(G)$ and considered the probability that the commutator of a randomly chosen pair of elements of $G$ equals a given element $g \in G$. We write $\Pr_g(G)$ to denote this probability. Motivated by their work, in this paper, we consider the probability that the autocommutator of a randomly chosen pair of elements, one from $G$ and the other from $\mathrm{Aut}(G)$, is equal to a given element $g \in G$. We write $\Pr_g(G, \mathrm{Aut}(G))$ to denote this probability. Thus,

$$\Pr_g(G, \mathrm{Aut}(G)) = \frac{|\{(x, \alpha) \in G \times \mathrm{Aut}(G) : [x, \alpha] = g\}|}{|G||\mathrm{Aut}(G)|}. \tag{1.2}$$

Notice that $\Pr_1(G, \mathrm{Aut}(G)) = \Pr(G, \mathrm{Aut}(G))$. Hence $\Pr_g(G, \mathrm{Aut}(G))$ is a generalization of $\Pr(G, \mathrm{Aut}(G))$. Clearly, $\Pr_g(G, \mathrm{Aut}(G)) = 1$ if and only if $K(G) = \{1\}$ and $g = 1$ if and only if $G = L(G)$ and $g = 1$. Also, $\Pr_g(G, \mathrm{Aut}(G)) = 0$ if and only if $g \notin S(G, \mathrm{Aut}(G))$. Therefore, we consider $G \neq L(G)$ and $g \in S(G, \mathrm{Aut}(G))$ throughout the paper.

In Section 2, we obtain a computing formula for $\Pr_g(G, \mathrm{Aut}(G))$ and deduce some of its consequences. In Section 3, we obtain several bounds for $\Pr_g(G, \mathrm{Aut}(G))$. In Section 4, we obtain some characterizations of $G$ through $\Pr(G, \mathrm{Aut}(G))$. Finally, in the last section, we show that $\Pr_g(G, \mathrm{Aut}(G))$ is an invariant under autoisoclinism of groups.

# 2 A computing formula

For any $x, g \in G$ let $T_{x,g}$ denotes the set $\{\alpha \in \mathrm{Aut}(G) : [x, \alpha] = g\}$. Note that $T_{x,1} = C_{\mathrm{Aut}(G)}(x)$. The following two lemmas play a crucial role in obtaining the computing formula for $\Pr_g(G, \mathrm{Aut}(G))$.

**Lemma 2.1.** *Let $G$ be a finite group. If $T_{x,g} \neq \phi$ then $T_{x,g} = \sigma C_{\mathrm{Aut}(G)}(x)$ for some $\sigma \in T_{x,g}$. Hence, $|T_{x,g}| = |C_{\mathrm{Aut}(G)}(x)|$.*

*Proof.* Let $\sigma \in T_{x,g}$ and $\beta \in \sigma C_{\mathrm{Aut}(G)}(x)$. Then $\beta = \sigma\alpha$ for some $\alpha \in C_{\mathrm{Aut}(G)}(x)$. We have

$$[x, \beta] = [x, \sigma\alpha] = x^{-1}\sigma(\alpha(x)) = [x, \sigma] = g.$$

Therefore, $\beta \in T_{x,g}$ and so $\sigma C_{\mathrm{Aut}(G)}(x) \subseteq T_{x,g}$. Again, let $\gamma \in T_{x,g}$ then $\gamma(x) = xg$. We have $\sigma^{-1}\gamma(x) = \sigma^{-1}(xg) = x$ and so $\sigma^{-1}\gamma \in C_{\mathrm{Aut}(G)}(x)$. Therefore, $\gamma \in \sigma C_{\mathrm{Aut}(G)}(x)$ which gives $T_{x,g} \subseteq \sigma C_{\mathrm{Aut}(G)}(x)$. Hence, the result follows. □

We know that $\mathrm{Aut}(G)$ acts on $G$ by the action $(\alpha, x) \mapsto \alpha(x)$ where $\alpha \in \mathrm{Aut}(G)$ and $x \in G$. Let $\mathrm{orb}(x) := \{\alpha(x) : \alpha \in \mathrm{Aut}(G)\}$ be the orbit of $x \in G$. Then by orbit-stabilizer theorem, we have

$$|\,\mathrm{orb}(x)| = \frac{|\,\mathrm{Aut}(G)|}{|C_{\mathrm{Aut}(G)}(x)|}. \tag{2.1}$$

**Lemma 2.2.** *Let $G$ be a finite group. Then $T_{x,g} \neq \phi$ if and only if $xg \in \mathrm{orb}(x)$.*

*Proof.* The result follows from the fact that $\alpha \in T_{x,g}$ if and only if $xg \in \mathrm{orb}(x)$. □

Now we state and prove the main theorem of this section.

**Theorem 2.3.** *Let $G$ be a finite group. If $g \in G$ then*

$$\Pr_g(G, \mathrm{Aut}(G)) = \frac{1}{|G||\,\mathrm{Aut}(G)|} \sum_{\substack{x \in G \\ xg \in \mathrm{orb}(x)}} |C_{\mathrm{Aut}(G)}(x)| = \frac{1}{|G|} \sum_{\substack{x \in G \\ xg \in \mathrm{orb}(x)}} \frac{1}{|\,\mathrm{orb}(x)|}.$$

*Proof.* We have $\{(x, \alpha) \in G \times \mathrm{Aut}(G) : [x, \alpha] = g\} = \bigsqcup_{x \in G}(\{x\} \times T_{x,g})$, where $\sqcup$ represents the union of disjoint sets. Therefore, by (1.2), we have

$$|G||\,\mathrm{Aut}(G)|\Pr_g(G, \mathrm{Aut}(G)) = |\bigsqcup_{x \in G}(\{x\} \times T_{x,g})| = \sum_{x \in G} |T_{x,g}|.$$

Hence, the result follows from Lemma 2.1, Lemma 2.2 and (2.1). □

Taking $g = 1$, in Theorem 2.3, we get the following corollary.

**Corollary 2.4.** *Let $G$ be a finite group. Then*

$$\Pr(G, \mathrm{Aut}(G)) = \frac{1}{|G||\,\mathrm{Aut}(G)|} \sum_{x \in G} |C_{\mathrm{Aut}(G)}(x)| = \frac{|\,\mathrm{orb}(G)|}{|G|}$$

*where* $\mathrm{orb}(G) = \{\mathrm{orb}(x) : x \in G\}$.

**Corollary 2.5.** *Let $G$ be a finite group. If $C_{\mathrm{Aut}(G)}(x) = \{I\}$ for all $x \in G \setminus \{1\}$, where $I$ is the identity element of* $\mathrm{Aut}(G)$*, then*

$$\Pr(G, \mathrm{Aut}(G)) = \frac{1}{|G|} + \frac{1}{|\,\mathrm{Aut}(G)|} - \frac{1}{|G||\,\mathrm{Aut}(G)|}.$$

*Proof.* By Corollary 2.4, we have

$$|G||\,\mathrm{Aut}(G)|\,\Pr(G, \mathrm{Aut}(G)) = \sum_{x \in G} |C_{\mathrm{Aut}(G)}(x)| = |\,\mathrm{Aut}(G)| + |G| - 1.$$

Hence, the result follows. □

We also have $|\{(x,\alpha) \in G \times \mathrm{Aut}(G) : [x,\alpha] = 1\}| = \sum_{\alpha \in \mathrm{Aut}(G)} |C_G(\alpha)|$ and hence

$$\Pr(G, \mathrm{Aut}(G)) = \frac{1}{|G||\mathrm{Aut}(G)|} \sum_{\alpha \in \mathrm{Aut}(G)} |C_G(\alpha)|. \tag{2.2}$$

We conclude this section with the following two results.

**Proposition 2.6.** *Let $G$ be a finite group. If $g \in G$ then*

$$\Pr_{g^{-1}}(G, \mathrm{Aut}(G)) = \Pr_g(G, \mathrm{Aut}(G)).$$

*Proof.* Let

$$A = \{(x,\alpha) \in G \times \mathrm{Aut}(G) : [x,\alpha] = g\} \text{ and}$$
$$B = \{(y,\beta) \in G \times \mathrm{Aut}(G) : [y,\beta] = g^{-1}\}.$$

Then $(x,\alpha) \mapsto (\alpha(x), \alpha^{-1})$ gives a bijection between $A$ and $B$. Therefore $|A| = |B|$. Hence the result follows from (1.2). □

**Proposition 2.7.** *Let $G$ and $H$ be two finite groups such that $\gcd(|G|,|H|) = 1$. If $(g,h) \in G \times H$ then*

$$\Pr_{(g,h)}(G \times H, \mathrm{Aut}(G \times H)) = \Pr_g(G, \mathrm{Aut}(G)) \Pr_h(H, \mathrm{Aut}(H)).$$

*Proof.* Let

$$\mathcal{X} = \{((x,y), \alpha_{G\times H}) \in (G \times H) \times \mathrm{Aut}(G \times H) : [(x,y), \alpha_{G\times H}] = (g,h)\},$$
$$\mathcal{Y} = \{(x, \alpha_G) \in G \times \mathrm{Aut}(G) : [x, \alpha_G] = g\} \text{ and}$$
$$\mathcal{Z} = \{(y, \alpha_H) \in H \times \mathrm{Aut}(H) : [y, \alpha_H] = h\}.$$

Since $\gcd(|G|,|H|) = 1$, by [1, Lemma 2.1], we have $\mathrm{Aut}(G \times H) = \mathrm{Aut}(G) \times \mathrm{Aut}(H)$. Therefore, for every $\alpha_{G\times H} \in \mathrm{Aut}(G \times H)$ there exist unique $\alpha_G \in \mathrm{Aut}(G)$ and $\alpha_H \in \mathrm{Aut}(H)$ such that $\alpha_{G\times H} = \alpha_G \times \alpha_H$, where $\alpha_G \times \alpha_H((x,y)) = (\alpha_G(x), \alpha_H(y))$ for all $(x,y) \in G \times H$. Also, for all $(x,y) \in G \times H$, we have $[(x,y), \alpha_{G\times H}] = (g,h)$ if and only if $[x,\alpha_G] = g$ and $[y,\alpha_H] = h$. These leads to show that $\mathcal{X} = \mathcal{Y} \times \mathcal{Z}$. Therefore

$$\frac{|\mathcal{X}|}{|G \times H||\mathrm{Aut}(G \times H)|} = \frac{|\mathcal{Y}|}{|G||\mathrm{Aut}(G)|} \cdot \frac{|\mathcal{Z}|}{|H||\mathrm{Aut}(H)|}.$$

Hence, the result follows from (1.1). □

# 3 Some bounds

We begin with the following lower bounds.

**Proposition 3.1.** *Let $G$ be a finite group. Then*

(a) $\Pr_g(G, \mathrm{Aut}(G)) \geq \frac{|L(G)|}{|G|} + \frac{|C_{\mathrm{Aut}(G)}(G)|(|G|-|L(G)|)}{|G||\mathrm{Aut}(G)|}$ *if $g = 1$.*

(b) $\Pr_g(G, \mathrm{Aut}(G)) \geq \frac{|L(G)||C_{\mathrm{Aut}(G)}(G)|}{|G||\mathrm{Aut}(G)|}$ *if $g \neq 1$.*

*Proof.* Let $\mathcal{C} := \{(x, \alpha) \in G \times \mathrm{Aut}(G) : [x, \alpha] = g\}$.

(a) We have $(L(G) \times \mathrm{Aut}(G)) \cup (G \times C_{\mathrm{Aut}(G)}(G)) \subseteq \mathcal{C}$ and $|(L(G) \times \mathrm{Aut}(G)) \cup (G \times C_{\mathrm{Aut}(G)}(G))|$ is equal to $|L(G)||\mathrm{Aut}(G)| + |C_{\mathrm{Aut}(G)}(G)||G| - |L(G)||C_{\mathrm{Aut}(G)}(G)|$. Hence, the result follows from (1.2).

(b) Since $g \in S(G, \mathrm{Aut}(G))$ we have $\mathcal{C}$ is non-empty. Let $(y, \beta) \in \mathcal{C}$ then $(y, \beta) \notin L(G) \times C_{\mathrm{Aut}(G)}(G)$ otherwise $[y, \beta] = 1$. It is easy to see that the coset $(y, \beta)(L(G) \times C_{\mathrm{Aut}(G)}(G))$ having order $|L(G)||C_{\mathrm{Aut}(G)}(G)|$ is a subset of $\mathcal{C}$. Hence, the result follows from (1.2). □

**Proposition 3.2.** *Let $G$ be a finite group. Then*

$$\mathrm{Pr}_g(G, \mathrm{Aut}(G)) \leq \mathrm{Pr}(G, \mathrm{Aut}(G)).$$

*The equality holds if and only if $g = 1$.*

*Proof.* By Theorem 2.3, we have

$$\begin{aligned}\mathrm{Pr}_g(G, \mathrm{Aut}(G)) &= \frac{1}{|G||\mathrm{Aut}(G)|} \sum_{\substack{x \in G \\ xg \in \mathrm{orb}(x)}} |C_{\mathrm{Aut}(G)}(x)| \\ &\leq \frac{1}{|G||\mathrm{Aut}(G)|} \sum_{x \in G} |C_{\mathrm{Aut}(G)}(x)| = \mathrm{Pr}(G, \mathrm{Aut}(G)).\end{aligned}$$

The equality holds if and only if $xg \in \mathrm{orb}(x)$ for all $x \in G$ if and only if $g = 1$. □

**Proposition 3.3.** *Let $G$ be a finite group and $p$ the smallest prime dividing $|\mathrm{Aut}(G)|$. If $g \neq 1$ then*

$$\mathrm{Pr}_g(G, \mathrm{Aut}(G)) \leq \frac{|G| - |L(G)|}{p|G|} < \frac{1}{p}.$$

*Proof.* By Theorem 2.3, we have

$$\mathrm{Pr}_g(G, \mathrm{Aut}(G)) = \frac{1}{|G|} \sum_{\substack{x \in G \setminus L(G) \\ xg \in \mathrm{orb}(x)}} \frac{1}{|\mathrm{orb}(x)|} \tag{3.1}$$

noting that for $x \in L(G)$ we have $xg \notin \mathrm{orb}(x)$. Also, for $x \in G \setminus L(G)$ and $xg \in \mathrm{orb}(x)$ we have $|\mathrm{orb}(x)| > 1$. Since $|\mathrm{orb}(x)|$ is a divisor of $|\mathrm{Aut}(G)|$ we have $|\mathrm{orb}(x)| \geq p$. Hence, the result follows from (3.1). □

The remaining part of this section is devoted in obtaining some lower and upper bounds for $\mathrm{Pr}(G, \mathrm{Aut}(G))$. The following theorem is an improvement of [8, Theorem 2.3 (ii)].

**Theorem 3.4.** *Let $G$ be a finite group and $p$ the smallest prime dividing $|\mathrm{Aut}(G)|$. Then*

$$\mathrm{Pr}(G, \mathrm{Aut}(G)) \geq \frac{|L(G)|}{|G|} + \frac{p(|G| - |X_G| - |L(G)|) + |X_G|}{|G||\mathrm{Aut}(G)|}$$

*and*

$$\mathrm{Pr}(G, \mathrm{Aut}(G)) \leq \frac{(p-1)|L(G)| + |G|}{p|G|} - \frac{|X_G|(|\mathrm{Aut}(G)| - p)}{p|G||\mathrm{Aut}(G)|},$$

*where $X_G = \{x \in G : C_{\mathrm{Aut}(G)}(x) = \{I\}\}$.*

*Proof.* We have $X_G \cap L(G) = \phi$. Therefore

$$\sum_{x\in G}|C_{\mathrm{Aut}(G)}(x)| = |X_G| + |\mathrm{Aut}(G)||L(G)| + \sum_{x\in G\setminus(X_G\cup L(G))}|C_{\mathrm{Aut}(G)}(x)|.$$

For $x \in G \setminus (X_G \cup L(G))$ we have $\{I\} \neq C_{\mathrm{Aut}(G)}(x) \neq \mathrm{Aut}(G)$ which implies $p \leq |C_{\mathrm{Aut}(G)}(x)| \leq \frac{|\mathrm{Aut}(G)|}{p}$. Therefore

$$\sum_{x\in G}|C_{\mathrm{Aut}(G)}(x)| \geq |X_G| + |\mathrm{Aut}(G)||L(G)| + p(|G| - |X_G| - |L(G)|) \tag{3.2}$$

and

$$\sum_{x\in G}|C_{\mathrm{Aut}(G)}(x)| \leq |X_G| + |\mathrm{Aut}(G)||L(G)| + \frac{|\mathrm{Aut}(G)|(|G| - |X_G| - |L(G)|)}{p}. \tag{3.3}$$

Hence, the result follows from Corollary 2.4, (3.2) and (3.3). □

We have the following two corollaries.

**Corollary 3.5.** *Let $G$ be a finite group. If $p$ and $q$ are the smallest primes dividing $|\mathrm{Aut}(G)|$ and $|G|$ respectively then*

$$\Pr(G, \mathrm{Aut}(G)) \leq \frac{p+q-1}{pq}.$$

*In particular, if $p = q$ then $\Pr(G, \mathrm{Aut}(G)) \leq \frac{2p-1}{p^2} \leq \frac{3}{4}$.*

*Proof.* Since $G \neq L(G)$ we have $|G : L(G)| \geq q$. Therefore, by Theorem 3.4, we have

$$\Pr(G, \mathrm{Aut}(G)) \leq \frac{1}{p}\left(\frac{p-1}{|G:L(G)|} + 1\right) \leq \frac{p+q-1}{pq}.$$

□

**Corollary 3.6.** *Let $G$ be a finite group and $p$, $q$ be the smallest primes dividing $|\mathrm{Aut}(G)|$ and $|G|$ respectively. If $G$ is non-abelian then*

$$\Pr(G, \mathrm{Aut}(G)) \leq \frac{q^2+p-1}{pq^2}.$$

*In particular, if $p = q$ then $\Pr(G, \mathrm{Aut}(G)) \leq \frac{p^2+p-1}{p^3} \leq \frac{5}{8}$.*

*Proof.* Since $G$ is non-abelian we have $|G : L(G)| \geq q^2$. Therefore, by Theorem 3.4, we have

$$\Pr(G, \mathrm{Aut}(G)) \leq \frac{1}{p}\left(\frac{p-1}{|G:L(G)|} + 1\right) \leq \frac{q^2+p-1}{pq^2}.$$

□

**Theorem 3.7.** *Let $G$ be a finite group. Then*

$$\Pr(G, \mathrm{Aut}(G)) \geq \frac{1}{|S(G, \mathrm{Aut}(G))|}\left(1 + \frac{|S(G, \mathrm{Aut}(G))| - 1}{|G : L(G)|}\right).$$

*The equality holds if and only if $\mathrm{orb}(x) = xS(G, \mathrm{Aut}(G))$ for all $x \in G \setminus L(G)$.*

*Proof.* For all $x \in G \setminus L(G)$ we have $\alpha(x) = x[x, \alpha] \in xS(G, \mathrm{Aut}(G))$. Therefore $\mathrm{orb}(x) \subseteq xS(G, \mathrm{Aut}(G))$ and so $|\mathrm{orb}(x)| \leq |S(G, \mathrm{Aut}(G))|$ for all $x \in G \setminus L(G)$. Now, by Corollary 2.4, we have

$$\begin{aligned}\Pr(G, \mathrm{Aut}(G)) &= \frac{1}{|G|}\left(\sum_{x \in L(G)} \frac{1}{|\mathrm{orb}(x)|} + \sum_{x \in G \setminus L(G)} \frac{1}{|\mathrm{orb}(x)|}\right) \\ &\geq \frac{|L(G)|}{|G|} + \frac{1}{|G|} \sum_{x \in G \setminus L(G)} \frac{1}{|S(G, \mathrm{Aut}(G))|}.\end{aligned}$$

Hence, the result follows. □

The following lemma is useful in obtaining the next corollary.

**Lemma 3.8.** *Let $G$ be a finite group. Then, for any two integers $m \geq n$, we have*

$$\frac{1}{n}\left(1 + \frac{n-1}{|G : L(G)|}\right) \geq \frac{1}{m}\left(1 + \frac{m-1}{|G : L(G)|}\right).$$

*If $L(G) \neq G$ then equality holds if and only if $m = n$.*

*Proof.* The proof is an easy exercise. □

**Corollary 3.9.** *Let $G$ be a finite group. Then*

$$\Pr(G, \mathrm{Aut}(G)) \geq \frac{1}{|K(G)|}\left(1 + \frac{|K(G)| - 1}{|G : L(G)|}\right).$$

*If $G \neq L(G)$ then the equality holds if and only if $K(G) = S(G, \mathrm{Aut}(G))$ and $\mathrm{orb}(x)$ is equal to $S(G, \mathrm{Aut}(G))$ for all $x \in G \setminus L(G)$.*

*Proof.* Since $|K(G)| \geq |S(G, \mathrm{Aut}(G))|$, the result follows from Theorem 3.7 and Lemma 3.8.

Note that the equality holds if and only if equality holds in Theorem 3.7 and Lemma 3.8. □

This lower bound is also obtained in [2]. Note that the lower bound in Theorem 3.7 is better than that in Corollary 3.9. Also

$$\frac{1}{|K(G)|}\left(1 + \frac{|K(G)| - 1}{|G : L(G)|}\right) \geq \frac{|L(G)|}{|G|} + \frac{p(|G| - |L(G)|)}{|G||\mathrm{Aut}(G)|}.$$

Hence, the lower bound in Theorem 3.7 is also better than that in [8, Theorem 2.3(ii)].

# 4 Some Characterizations

In [2], Arora and Karan obtain a characterization of a finite group $G$ if equality holds in Corollary 3.9. In this section, we obtain some characterizations of $G$ if equality holds in Corollary 3.5 and Corollary 3.6. We begin with the following result.

**Proposition 4.1.** *Let $G$ be a finite group with $\Pr(G, \mathrm{Aut}(G)) = \frac{p+q-1}{pq}$ for some primes $p$ and $q$. Then $pq$ divides $|G||\mathrm{Aut}(G)|$. If $p$ and $q$ are the smallest primes dividing $|\mathrm{Aut}(G)|$ and $|G|$ respectively, then*

$$\frac{G}{L(G)} \cong \mathbb{Z}_q.$$

*In particular, if $G$ and $\mathrm{Aut}(G)$ are of even order and $\Pr(G, \mathrm{Aut}(G)) = \frac{3}{4}$ then $\frac{G}{L(G)} \cong \mathbb{Z}_2$.*

*Proof.* By (1.1), we have $(p+q-1)|G||\mathrm{Aut}(G)| = pq|\{(x,\alpha) \in G \times \mathrm{Aut}(G) : [x,\alpha] = 1\}|$. Therefore, $pq$ divides $|G||\mathrm{Aut}(G)|$.

If $p$ and $q$ are the smallest primes dividing $|\mathrm{Aut}(G)|$ and $|G|$ respectively then, by Theorem 3.4, we have

$$\frac{p+q-1}{pq} \leq \frac{1}{p}\left(\frac{p-1}{|G : L(G)|} + 1\right)$$

which gives $|G : L(G)| \leq q$. Hence, $\frac{G}{L(G)} \cong \mathbb{Z}_q$. □

**Proposition 4.2.** *Let $G$ be a finite non-abelian group with $\Pr(G, \mathrm{Aut}(G)) = \frac{q^2+p-1}{pq^2}$ for some primes $p$ and $q$. Then $pq$ divides $|G||\mathrm{Aut}(G)|$. If $p$ and $q$ are the smallest primes dividing $|\mathrm{Aut}(G)|$ and $|G|$ respectively then*

$$\frac{G}{L(G)} \cong \mathbb{Z}_q \times \mathbb{Z}_q.$$

*In particular, if $G$ and $\mathrm{Aut}(G)$ are of even order and $\Pr(G, \mathrm{Aut}(G)) = \frac{5}{8}$ then $\frac{G}{L(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.*

*Proof.* By (1.1), we have $(q^2+p-1)|G||\mathrm{Aut}(G)| = pq^2|\{(x,\alpha) \in G \times \mathrm{Aut}(G) : [x,\alpha] = 1\}|$. Therefore, $pq$ divides $|G||\mathrm{Aut}(G)|$.

If $p$ and $q$ are the smallest primes dividing $|\mathrm{Aut}(G)|$ and $|G|$ respectively then, by Theorem 3.4, we have

$$\frac{q^2+p-1}{pq^2} \leq \frac{1}{p}\left(\frac{p-1}{|G : L(G)|} + 1\right)$$

which gives $|G : L(G)| \leq q^2$. Since $G$ is non-abelian we have $|G : L(G)| \neq 1, q$. Hence, $\frac{G}{L(G)} \cong \mathbb{Z}_q \times \mathbb{Z}_q$. □

We conclude this section with the following partial converses of Proposition 4.1 and 4.2.

**Proposition 4.3.** *Let $G$ be a finite group. Let $p, q$ be the smallest prime divisors of $|\mathrm{Aut}(G)|$ and $|G|$ respectively and $|\mathrm{Aut}(G) : C_{\mathrm{Aut}(G)}(x)| = p$ for all $x \in G \setminus L(G)$.*

(a) *If $\frac{G}{L(G)} \cong \mathbb{Z}_q$ then $\Pr(G, \mathrm{Aut}(G)) = \frac{p+q-1}{pq}$.*

(b) *If $\frac{G}{L(G)} \cong \mathbb{Z}_q \times \mathbb{Z}_q$ then $\Pr(G, \mathrm{Aut}(G)) = \frac{q^2+p-1}{pq^2}$.*

*Proof.* Since $|\operatorname{Aut}(G) : C_{\operatorname{Aut}(G)}(x)| = p$ for all $x \in G \setminus L(G)$ we have $|C_{\operatorname{Aut}(G)}(x)| = \frac{|\operatorname{Aut}(G)|}{p}$ for all $x \in G \setminus L(G)$. Therefore, by Corollary 2.4, we have

$$\begin{aligned}\Pr(G, \operatorname{Aut}(G)) &= \frac{|L(G)|}{|G|} + \frac{1}{|G||\operatorname{Aut}(G)|} \sum_{x \in G \setminus L(G)} |C_{\operatorname{Aut}(G)}(x)| \\ &= \frac{|L(G)|}{|G|} + \frac{|G| - |L(G)|}{p|G|}.\end{aligned}$$

Thus

$$\Pr(G, \operatorname{Aut}(G)) = \frac{1}{p}\left(\frac{p-1}{|G : L(G)|} + 1\right). \tag{4.1}$$

(a) If $\frac{G}{L(G)} \cong \mathbb{Z}_q$ then (4.1) gives $\Pr(G, \operatorname{Aut}(G)) = \frac{p+q-1}{pq}$.

(b) If $\frac{G}{L(G)} \cong \mathbb{Z}_q \times \mathbb{Z}_q$ then (4.1) gives $\Pr(G, \operatorname{Aut}(G)) = \frac{q^2+p-1}{pq^2}$. □

# 5 Autoisoclinism of groups

In the year 1940, Hall defined isoclinism between two groups (see [5]). In the year 1995, Lescot [7] showed that the commuting probability of two isoclinic finite groups are same. Later on Pournaki and Sobhani [10] showed that $\Pr_g(G) = \Pr_{\beta(g)}(H)$ if $(\alpha, \beta)$ is an isoclinism between the finite groups $G$ and $H$. Following Hall, Moghaddam et al. [9] have defined autoisoclinism between two groups. Recall that two groups $G$ and $H$ are said to be autoisoclinic if there exist isomorphisms $\psi : \frac{G}{L(G)} \to \frac{H}{L(H)}$, $\beta : K(G) \to K(H)$ and $\gamma : \operatorname{Aut}(G) \to \operatorname{Aut}(H)$ such that the following diagram commutes

$$\begin{array}{ccc} \frac{G}{L(G)} \times \operatorname{Aut}(G) & \xrightarrow{\psi \times \gamma} & \frac{H}{L(H)} \times \operatorname{Aut}(H) \\ \downarrow{\scriptstyle a_{(G,\operatorname{Aut}(G))}} & & \downarrow{\scriptstyle a_{(H,\operatorname{Aut}(H))}} \\ K(G) & \xrightarrow{\beta} & K(H) \end{array}$$

where the maps $a_{(G,\operatorname{Aut}(G))} : \frac{G}{L(G)} \times \operatorname{Aut}(G) \to K(G)$ and $a_{(H,\operatorname{Aut}(H))} : \frac{H}{L(H)} \times \operatorname{Aut}(H) \to K(H)$ are given by

$$a_{(G,\operatorname{Aut}(G))}(xL(G), \alpha_1) = [x, \alpha_1] \text{ and } a_{(H,\operatorname{Aut}(H))}(yL(H), \alpha_2) = [y, \alpha_2]$$

respectively. In this case, the pair $(\psi \times \gamma, \beta)$ is called an autoisoclinism between the groups $G$ and $H$.

Recently, in [12], Rismanchain and Sepehrizadeh have shown that $\Pr(G, \operatorname{Aut}(G)) = \Pr(H, \operatorname{Aut}(H))$ if $G$ and $H$ are autoisoclinic finite groups. We conclude this paper with the following generalization of [12, Lemma 2.5].

**Theorem 5.1.** *Let $G$ and $H$ be two finite groups and $(\psi \times \gamma, \beta)$ an autoisoclinism between them. Then*

$$\Pr_g(G, \operatorname{Aut}(G)) = \Pr_{\beta(g)}(H, \operatorname{Aut}(H)).$$

*Proof.* Let $\mathcal{S}_g = \{(xL(G), \alpha_1) \in \frac{G}{L(G)} \times \operatorname{Aut}(G) : [x, \alpha_1] = g\}$ and $\mathcal{T}_{\beta(g)} = \{(yL(H), \alpha_2) \in \frac{H}{L(H)} \times \operatorname{Aut}(H) : [y, \alpha_2] = \beta(g)\}$. Since $(\psi \times \gamma, \beta)$ an autoisoclinism between $G$ and

$H$, the mapping $\lambda : \mathcal{S}_g \to \mathcal{T}_{\beta(g)}$ given by $(xL(G), \alpha_1) \mapsto (\psi(xL(G)), \gamma(\alpha_1))$ gives an one to one correspondence between $\mathcal{S}_g$ and $\mathcal{T}_{\beta(g)}$. Hence, $|\mathcal{S}_g| = |\mathcal{T}_{\beta(g)}|$. Also

$$|\{(x, \alpha_1) \in G \times \mathrm{Aut}(G) : [x, \alpha_1] = g\}| = |L(G)||\mathcal{S}_g| \tag{5.1}$$

and

$$|\{(y, \alpha_2) \in H \times \mathrm{Aut}(H) : [y, \alpha_2] = \beta(g)\}| = |L(H)||\mathcal{T}_{\beta(g)}|. \tag{5.2}$$

Therefore, by (1.1) and (5.1), we have

$$\mathrm{Pr}_g(G, \mathrm{Aut}(G)) = \frac{|L(G)||\mathcal{S}_g|}{|G||\,\mathrm{Aut}(G)|}.$$

Since $|G/L(G)| = |H/L(H)|$ and $|\,\mathrm{Aut}(G)| = |\,\mathrm{Aut}(H)|$ we have

$$\mathrm{Pr}_g(G, \mathrm{Aut}(G)) = \frac{|L(H)||\mathcal{T}_{\beta(g)}|}{|H||\,\mathrm{Aut}(H)|}.$$

Hence, the result follows from (5.2) and (1.1). □

# References


[1] J. H. Christopher and L. R. Darren, *Automorphism of finite abelian groups*, Amer. Math. Monthly, **114**, 917–923.

[2] H. Arora and R. Karan, *What is the probability an automorphism fixes a group element?*, Comm. Algebra, **45**(3), 1141–1150 (2017).

[3] A. K. Das, R. K. Nath and M. R. Pournaki, *A survey on the estimation of commutativity in finite groups*, Southeast Asian Bull. Math., **37**(2), 161–180 (2013).

[4] P. Erdös and P. Turán, *On some problems of a statistical group-theory. IV*, Acta. Math. Acad. Sci. Hungar., **19**, 413–435 (1968).

[5] P. Hall, *The classification of prime power groups*, J. Reine Angew. Math., **182**, 130–141 (1940).

[6] P. V. Hegarty, *The absolute centre of a group*, J. Algebra, **169**(3), 929–935 (1994).

[7] P. Lescot, *Isoclinism classes and commutativity degrees of finite groups*, J. Algebra, **177**, 847–869 (1995).

[8] M. R. R. Moghaddam, F. Saeedi and E. Khamseh, *The probability of an automorphism fixing a subgroup element of a finite group*, Asian-Eur. J. Math. **4**(2), 301308 (2011).

[9] M. R. R. Moghaddam, M. J. Sadeghifard and M. Eshrati, *Some properties of autoisoclinism of groups*, Fifth International group theory conference, Islamic Azad University, Mashhad, Iran, 13-15 March 2013.

[10] M. R. Pournaki and R. Sobhani, *Probability that the commutator of two group elements is equal to a given element*, *J. Pure Appl. Algebra*, **212**, 727–734 (2008).

[11] R. Rezaei, A. Erfanian and P. Lescot, *A note on the relative commutativity degree of finite groups*, Asian-Eur. J. Math., **7** (2014), DOI: 10.1142/S179355711450017X.

[12] M. R. Rismanchian and Z. Sepehrizadeh, *Autoisoclinism classes and autocommutativity degrees of finite groups*, Hacet. J. Math. Stat. **44**(4), 893–899 (2015).

[13] G. J. Sherman, *What is the probability an automorphism fixes a group element?*, Amer. Math. Monthly, **82**, 261–264 (1975).